arXiv:1609.01062v1 [math.CV] 5 Sep 2016

CLASSIFICATION RESULTS OF TOTALLY REAL IMMERSIONS AND EMBEDDINGS INTO $\mathbb{C}^N$

MARKO SLAPAR AND RAFAEL TORRES

Abstract: We prove that the existence of totally real immersions of manifolds is a closed property under cut-and-paste constructions along submanifolds including connected sums. We study the existence of totally real embeddings for simply connected 5-manifolds and orientable 6-manifolds and determine the diffeomorphism and homotopy types. We show that high-dimensional manifolds with prescribed fundamental group admit a totally real embedding in contrast with the situation in dimension four.

1. Introduction and main results

In this paper, we are interested in the following kind of maps.

Definition 1. Let M^N be a closed smooth N-manifold and let J denote the standard complex structure on the tangent bundle of $\mathbb{C}^N$. An immersion $M^N \to \mathbb{C}^N$ is totally real if if the tangent bundle TM^N contains no complex lines, i.e., if

$$TM^N \cap JTM^N = \{0\} \tag{1.1}$$

at all points of TM^N. An embedding $M^N \hookrightarrow \mathbb{C}^N$ that satisfies (1.1) is called a totally real embedding.

A canonical problem is to distinguish between those manifolds that admit totally real immersions and the ones who admit totally real embeddings. This dichotomy is already somewhat interesting in the case of manifolds with simple topology. For example, every N-sphere S^N admits a totally real immersion into $\mathbb{C}^N$, yet no totally real embedding exists if $N > 3$; see Gromov [8, p. 193], Stout-Zame [22]. We occupy ourselves with the study of the distinction of the maps of Definition 1 for a large class of manifolds in this paper. Necessary and sufficient topological conditions for the existence of totally real immersions and embeddings have been studied by Gromov [7], Wells [24], Audin [2], Forstnerič [5], Gong [6] and Jacobowitz-Landweber [10] among several other mathematicians (see Section 2).

Our first two theorems state that the existence of totally real immersions is a property which is closed under certain fundamental cut-and-paste constructions of manifolds along submanifolds.

Theorem A. *Let M^N be a closed smooth N-manifold that admits a totally real immersion into $\mathbb{C}^N$ and let $i : S^p \times D^{N-p} \hookrightarrow M^N$ be a smooth embedding for $0 \leq p \leq 2$. The N-manifold*

$$\hat{M}^N := M^N \backslash i(S^p \times D^{N-p}) \cup (D^{p+1} \times S^{N-p-1}) \tag{1.2}$$

that is obtained by performing surgery along $i(S^p \times \{0\})$ admits a totally real immersion $\hat{M}^N \to \mathbb{C}^N$.

2010 *Mathematics Subject Classification.* 57R40, 3240, 32Q99.

Theorem B. *Let M_1^N and M_2^N be closed smooth N-manifolds that admit a totally real immersion into $\mathbb{C}^N$. There is a totally real immersion $M_1^N \# M_2^N \to \mathbb{C}^N$.*

A circle has a totally real embedding into $\mathbb{C}$ and every closed orientable surface admits a totally real immersion into $\mathbb{C}^2$, yet the 2-torus is the only orientable closed surface that admits a totally real embedding. Ahern-Rudin [1] constructed an explicit totally real embedding $S^3 \hookrightarrow \mathbb{C}^3$ and Forstnerič showed that every closed orientable 3-manifold admits a totally real embedding into $\mathbb{C}^3$ [5, 1.4 Theorem]. Jacobowitz-Landweber have shown that a necessary and sufficient condition for a closed smooth orientable 4-manifold M^4 to admit a totally real immersion into $\mathbb{C}^4$ is the vanishing of its first Pontrjagin class $p_1(M^4)$ [11, Corollary 4.1]. Our next result address the situation in dimension five using the classification of closed simply connected 5-manifolds of Barden [3] and Smale [21]. A smooth manifold M^N is irreducible if for every connected sum decomposition $M^N = M_1 \# M_2$, either M_1 or M_2 is diffeomorphic to the n-sphere. The nontrivial 3-sphere bundle over the 2-sphere is denoted by $S^3 \widetilde{\times} S^2$.

Theorem C. *Every closed smooth simply connected 5-manifold M^5 admits a totally real immersion*

$$M^5 \to \mathbb{C}^5. \tag{1.3}$$

Let M^5 be an irreducible simply connected 5-manifold. There is a totally real embedding

$$M^5 \hookrightarrow \mathbb{C}^5 \tag{1.4}$$

if and only if

$$M^5 \in \{\mathrm{SU}(3)/\mathrm{SO}(3), S^2 \times S^3, S^3 \widetilde{\times} S^2\} \tag{1.5}$$

up to diffeomorphism.

The 5-sphere is the only irreducible simply connected 5-manifold that does not admit a totally real embedding into complex 5-space. A complete list of simply connected 5-manifolds that admit such an embedding is given in Section 3.2 and it includes the following set of examples.

Corollary D. *Let M^5 be a closed smooth simply connected 5-manifold with torsion-free second homology group $H_2(M^5;\mathbb{Z})$ and suppose $k \in \mathbb{N}$. There is a totally real embedding*

$$M^5 \hookrightarrow \mathbb{C}^5 \tag{1.6}$$

if and only if M^5 is diffeomorphic to

$$S^3 \widetilde{\times} S^2 \# (2k-2)(S^2 \times S^3) \tag{1.7}$$

if $\omega_2(M^5) \neq 0$ and to

$$(2k-1)(S^2 \times S^3) \tag{1.8}$$

otherwise.

The homotopy type of a closed simply connected 5-manifold determines its diffeomorphism class [3, Section 2]. This is no longer the case in dimension six, where closed homotopy equivalent 6-manifolds need not be homeomorphic. Building on results of Wall [23], our next theorem states in terms of characteristic classes, the necessary and sufficient conditions for the existence of the maps of Definition 1

for 6-manifolds. The symbol $\chi(M^N)$ stands for the Euler characteristic of the N-manifold M^N.

Theorem E. *Let M^6 be a closed smooth orientable 6-manifold without 2-torsion in $H^3(M^6;\mathbb{Z})$. There is a totally real immersion*

$$M^6 \to \mathbb{C}^6 \tag{1.9}$$

if and only if the first Pontrjagin class satisfies $p_1(M^6) = 0$.

There is a totally real embedding

$$M^6 \hookrightarrow \mathbb{C}^6 \tag{1.10}$$

if and only if

$$p_1(M^6) = 0 = \chi(M^6). \tag{1.11}$$

Results of Dehn [4], Kervaire-Milnor [18] and Gromov [7] imply that there is a totally real immersion $M^N(G) \to \mathbb{C}^N$ of a closed orientable N-manifold $M^N(G)$ with prescribed finitely presented fundamental group for every $N \geq 4$. The fundamental group does impose a restriction for the existence of totally real embeddings into $\mathbb{C}^4$ since an argument due to Wells [24] shows that the Euler characteristic of such a 4-manifold must be zero. Our next result shows that this is not the case in high dimensions.

Theorem F. *Let G be a finitely presented group. There is a totally real immersion*

$$M^4(G) \to \mathbb{C}^4 \tag{1.12}$$

of a closed smooth orientable 4-manifold $M^4(G)$ with fundamental group isomorphic to G.

There is a totally real embedding

$$M^N(G) \hookrightarrow \mathbb{C}^N, \tag{1.13}$$

where $M^N(G)$ is a closed smooth N-manifold with fundamental group isomorphic to G and for every $N \geq 5$.

The structure of the paper is as follows. Classical and new existence results on totally real immersions and embeddings which we build upon to prove the results presented in the introduction are collected in Section 2. A proof of Theorem A and Theorem B is given in Section 3.1. Theorem C and its corollary are proven in Section 3.2 and Section 3.3, respectively, while Theorem E is proven in Section 3.4 and Theorem F in Section 3.5. Section 4 contains some results addressing the existence of more general immersions with respect to the ones considered in Definition 1 (cf. [10]).

2. Totally real immersions and embeddings

Wells showed that if an N-manifold M^N admits a totally real immersion into $\mathbb{C}^N$ in the sense of Definition 1, then its complexified tangent bundle

$$\mathbb{C}TM^N := TM^N \otimes \mathbb{C} \tag{2.1}$$

is trivial [24]. Gromov proved that if $\mathbb{C}TM$ is trivial, then such an immersion exists by using convex integration in [7] (cf. [10, Theorem 1.2]).

Theorem 1. *Wells* [24]*, Gromov* [7]*. There is a totally real immersion $M^N \to \mathbb{C}^N$ if and only if $\mathbb{C}TM^N$ is a trivial bundle of rank N.*

Let ϵ^k be a trivial rank k bundle over M^N. If the Whitney sum $TM^N \oplus \epsilon^1$ is a trivial bundle, then so is its complexification $\mathbb{C}TM^N$. The following result is an immediate consequence of the work of Gromov

Corollary 1. *Every stably parallelizable manifold M^N admits a totally real immersion into $\mathbb{C}^N$.*

Audin studied conditions under which the product of manifolds admits a totally real immersion whenever each of the factors does [2, 6.2].

Proposition 1. *Audin* [2, 6.2.3. Remarque]. *Suppose there exist a totally real immersion $X^N \to \mathbb{C}^N$ and a totally real embedding $Y^M \to \mathbb{C}^M$. There is a totally real embedding*

$$X^N \times Y^M \hookrightarrow \mathbb{C}^N \times \mathbb{C}^M \cong \mathbb{C}^{N+M}. \tag{2.2}$$

Our next result points extends her results to the more general case of fiber bundles.

Proposition 2. *Let X^N be a closed smooth orientable N-manifold with trivial complexified tangent bundle $\mathbb{C}TX^N$. Let Y^{N+k} be the total space of a k-torus bundle for $k \in \mathbb{N}$*

$$T^k \hookrightarrow Y^{N+k} \xrightarrow{\pi} X^N. \tag{2.3}$$

There exists a totally real immersion

$$Y^{N+k} \to \mathbb{C}^{N+k}. \tag{2.4}$$

Notice that the converse to the conclusion of Proposition 2 does not hold as exemplified by $S^1 \hookrightarrow S^5 \to \mathbb{CP}^2$. Moreover, the conclusion of Proposition 2 can be strengthened to cover embeddings.

Proof. The tangent bundle of Y^{N+k} can be written as

$$TY^{N+k} \cong \pi^* TX^N \oplus \epsilon^k \tag{2.5}$$

where ϵ^k is a trivial rank k bundle over Y^{N+k} whose elements are tangent to the torus fibers [20]. Its complexification is

$$\mathbb{C}TY^{N+k} = TY^{N+k} \otimes \mathbb{C} = (\pi^* TX^N \oplus \epsilon^k) \otimes \mathbb{C} = (\pi^* TX^N \otimes \mathbb{C}) \oplus (Y^{N+k} \times \mathbb{C}), \tag{2.6}$$

which is trivial since $\mathbb{C}TX^N$ is assumed to be trivial. Theorem 1 implies that there is a totally real immersion $Y^{N+k} \to \mathbb{C}^{N+k}$. □

Recall that the Kervaire semi-characteristic of a closed smooth n-manifold M^N of dimension $N = 2k+1$ for $k \in \mathbb{N}$ is defined as

$$\hat{\chi}_{\mathbb{Z}/2}(M^N) := \sum_{i=0}^{k} \dim H^i(M^N; \mathbb{Z}/2) \bmod 2 \tag{2.7}$$

by Luzstig-Milnor-Peterson [17].

Theorem 2. *Audin* [2, 0.4 Proposition, 0.5 Théorème, 0.6 Corollaire]. *Let M^N be a closed smooth connected orientable N-manifold of dimension and suppose there is a totally real immersion $M^N \to \mathbb{C}^N$.*

(A) Suppose N is even. There is a totally real embedding $M^N \hookrightarrow \mathbb{C}^N$ if and only if $\chi(M^N) = 0$.

(B) Suppose $N = 4k+1$ for $k \in \mathbb{N}$. There is a totally real embedding $M^N \hookrightarrow \mathbb{C}^N$ if and only if $\hat{\chi}_{\mathbb{Z}/2}(M^N) = 0$.

We finish the section with the following sets of examples.

Proposition 3. *Let M^5 be a closed smooth simply connected 5-manifold with second Stiefel-Whitney class $\omega_2(M^5) = 0$.*

The manifold M^5 is stably parallelizable and there is a totally real immersion $M^5 \to \mathbb{C}^5$.

There is a totally real embedding $M^5 \hookrightarrow \mathbb{C}^5$ if and only if M^5 is parallelizable.

Proof. A result of Hirsch states that an n-manifold M is stably parallelizable if and only if M is orientable and it immerses into $\mathbb{R}^{n+1}$ [9] (cf. [8, Section 1.1.3]). Barden has shown a closed simply connected 5-manifold admits an immersion into $\mathbb{R}^6$ if and only if its second Stiefel-Whitney class vanishes [3, Lemma 2.4]. The existence of the totally real immersion now follows from Corollary 1.

Item (B) of Theorem 2 implies that there exists a totally real embedding of M^5 into $\mathbb{C}^5$ if and only if the Kervaire semi-characteristic $\hat{\chi}_{\mathbb{Z}/2}(M^5) = 0$. Kervaire has shown in [13] that the only obstruction for a stably parallelizable odd-dimensional manifold to be parallelizable, is the vanishing of the Kervaire semi-characteristic provided that the dimension is not one, three, nor seven.

□

Smale's [21] and Barden's [3] classification of closed simply connected 5-manifolds implies that a manifold of Proposition 3 is diffeomorphic to a connected sum

$$S^5 \# (k-1)(S^2 \times S^3) \# (k_1 - 1) M_{p_1^k} \# \cdots \# (k_i - 1) M_{p_i^k} \tag{2.8}$$

for $k, k_1, \ldots, k_j \in \mathbb{N}$, where each manifold $M_{p_i^k}$ has $H^2(M_{p_i^k}; \mathbb{Z}) = \mathbb{Z}/p_i^k \oplus \mathbb{Z}/p_i^k$ and $\omega_2(M_{p_i^k}) = 0$; see Table 1. In particular, the manifold (2.8) admits a totally real embedding into complex 5-space if and only if k is a positive even number.

Appealing to another classification result due to Smale [21] we obtain the following six-dimensional examples.

Proposition 4. *Every closed smooth 2-connected 6-manifold M^6 admits a totally real immersion*

$$M^6 \to \mathbb{C}^6 \tag{2.9}$$

and there is a totally real embedding

$$M^6 \hookrightarrow \mathbb{C}^6 \tag{2.10}$$

if and only if M^6 is diffeomorphic to $S^3 \times S^3$.

Proof. Smale [21] has shown that a closed smooth 2-connected 6-manifold M is diffeomorphic to S^6 or to a connected sum $n(S^3 \times S^3)$ of n copies of the product of two 3-spheres for $n \in \mathbb{N}$. Since the connected sum of two stably parallelizable manifolds is stably parallelizable, it follows that every closed smooth 2-connected 6-manifold is stably parallelizable. Corollary 1 implies that the totally real immersion (2.9) exists for every such 6-manifold. Theorem 2 and Smale's cited classification result imply that there is a totally real embedding (2.10) if and only if M^6 is diffeomorphic to $S^3 \times S^3$. We point out that Ahern-Rudin have given an explicit construction of a totally real embedding $S^3 \hookrightarrow \mathbb{C}^3$ [1] that can be used to construct a totally real embedding of the product of two 3-spheres into $\mathbb{C}^6$ (cf. [2, 6.2.3 Remarque]).

□

3. Proofs

3.1. **Proof of Theorems A and Theorem B.** We first show that the manifold

$$\hat{M}^N := M^N \backslash i(S^2 \times D^{N-2}) \cup (D^3 \times S^{N-3}) \tag{3.1}$$

that is obtained by performing surgery along $i(S^2 \times \{0\})$ admits a totally real immersion $\hat{M}^N \to \mathbb{C}^N$ for clarity purposes, and then discuss the corresponding generalization to the cases $p = 0, 1$. Theorem 1 states that we need to show that $\mathbb{C}T\hat{M}^N$ is trivial. Set $S := i(S^2 \times \{0\})$ and let $\{e_1, e_2, e_3, \dots, e_N\}$ be sections that trivialize the bundle $\mathbb{C}TM^N$. We can assume that along the 2-sphere S, the elements $\{e_1, e_2\}$ give a trivialization of $\mathbb{C}TS$ and the elements $\{e_3, \dots, e_N\}$ trivialize the normal bundle NS of S. This is justified by the following argument. Let $\{f_1, f_2, f_3, \dots, f_N\}$ be nowhere zero sections of $\mathbb{C}TM^N|_S$, defined over S, so that $\{f_1, f_2\}$ trivialize $\mathbb{C}TS$ and $\{f_3, \dots, f_N\}$ trivialize NS. There exists a map

$$A : S \to \mathrm{GL}(N, \mathbb{C}), \tag{3.2}$$

so that $f_i = Ae_i$ for $i = 1, 2, \dots, n$. Since $\pi_2(\mathrm{GL}(N, \mathbb{C})) = 0$, there is a homotopy A_t, so that $A_t = A$ near $t = 0$ and $A_t = I$ near $t = 1$. We can use this homotopy to connect the trivialization $\{f_1, f_2, f_3, \dots, f_N\}$ over S with the trivialization $\{e_1, e_2, e_3, \dots, e_N\}$ outside a neighborhood of S.

Let now W be the standard cobordism between M^N and $\hat{M}^N$ that is obtained by attaching an $N+1$-dimensional 3-handle $H := D^3 \times D^{N-2}$ to $M^N \times [0, 1]$ along $M^N \times \{1\}$ via the gluing map $i_{S^2 \times D^{N-2}} \times \{1\}$. We proceed to show that $\mathbb{C}TW$ is a trivial bundle. Let e be inward normal vector field to the boundary sphere $S = S^2 \times \{0\}$ in the core $D := D^3 \times \{0\}$ of the handle. Since $\pi_2(\mathrm{GL}(3, \mathbb{C})) = 0$, we can extend $\{e, e_1, e_3\}$ to a trivialization of $\mathbb{C}TD$. We can also trivially extend $\{e_3, \dots, e_N\}$ to a nonzero trivialization of the normal bundle of D in the handle H; recall $\pi_2(\mathrm{O}(N-2)) = 0$ and there are no framing issues attaching 3-handles. Since $M \cup D$ is a deformation retract of W, $\{e, e_1, e_2, e_3, \dots, e_N\}$ can be extended to give a trivialization of $\mathbb{C}TW$. Since along $\hat{M}^N = \partial_+ W$, the tangent bundle TW is a Whitney sum $T\hat{M}^N \oplus \epsilon^1$ with a trivial line bundle ϵ^1, we conclude that $\mathbb{C}T\hat{M}^N \oplus (\epsilon^1 \otimes \mathbb{C})$ is trivial. Since $\mathbb{C}T\hat{M}^N$ is stably trivial, then it is a trivial bundle [10, Lemma 1.2].

Remark 1. *Small tweaks to the proof of Theorem A yield the same conclusion for surgeries performed to M^N along*

$$S^p \times D^{n-p} \hookrightarrow M^n \tag{3.3}$$

for $p = 0, 1$ whose normal bundle over the handle is trivial. In these cases there are two choices of framings $\pi_0(\mathrm{O}(N)) = \pi_1(\mathrm{O}(N-1)) = \mathbb{Z}/2$ (except for $N = 3$ when $\pi_1(\mathrm{O}(2)) = \mathbb{Z}$). For exactly one choice of framing, we can extend the trivialisation of the normal bundle from the boundary of the core of the handle to its core as in the previous proof.

The above remark in the case $p = 0$ yields Theorem B as a corollary.

3.2. **Proof of Theorem C.** The case of 5-manifolds with vanishing second Stiefel-Whitney class was settled in Proposition 3. The classification results of closed simply connected 5-manifolds up to diffeomorphism of Barden [3] and Smale [21] imply that any such 5-manifold is difeomorphic to a connected sum of manifolds

5-manifold Y	$H^2(Y;\mathbb{Z})$	$\omega_2(Y)$	$\hat{\chi}_{\mathbb{Z}/2}(Y)$
S^5	0	0	1
$S^2 \times S^3$	$\mathbb{Z}$	0	0
M_{p^k}	$\mathbb{Z}/p^k \oplus \mathbb{Z}/p^k$	0	1
$\mathrm{SU}(3)/\mathrm{SO}(3)$	$\mathbb{Z}/2$	$\neq 0$	0
$S^3\widetilde{\times}S^2$	$\mathbb{Z}$	$\neq 0$	0
X_k	$\mathbb{Z}/2^k \oplus \mathbb{Z}/2^k$	$\neq 0$	1

TABLE 1. Building blocks of simply connected 5-manifolds ($k \in \mathbb{N}$)

in Table 1. We proceed to argue that every manifold in the table has trivial complexified tangent bundle and then invoke Theorem B and Theorem 1 in order to prove the first part of Theorem C. Therefore, we need to show that the Wu manifold $\mathrm{SU}(3)/\mathrm{SO}(3)$, the nontrivial bundle $S^3\widetilde{\times}S^2$, and the manifold X_k with k have trivial complexified tangent bundle. Audin has shown that the Wu manifold admits a totally real embedding into $\mathbb{C}^5$ [2, Proposition 0.8]. The same conclusion holds for $S^3\widetilde{\times}S^2$ by Proposition 2 since it is the total space of a circle bundle

$$S^1 \hookrightarrow S^3\widetilde{\times}S^2 \to \mathbb{CP}^2\#\overline{\mathbb{CP}^2}, \tag{3.4}$$

where the base 4-manifold has trivial complexified tangent bundle. We now show that $\mathbb{C}TX_k$ is a trivial bundle for every value $k \in \mathbb{N}$. Let $S \hookrightarrow S^3\widetilde{\times}S^2$ be a 2-sphere that represents 2^k-times the generator of the infinite cyclic group $H_2(S^3\widetilde{\times}S^2;\mathbb{Z})$. Notice that the normal bundle of S is trivial. The manifold X_k is obtained from $S^3\widetilde{\times}S^2$ by performing the surgery of Theorem A along S and therefore $\mathbb{C}TX_k$ is trivial. The claim regarding totally real embeddings into complex 5-space follows immediately from Theorem 2.

□

Remark 2. *Every closed simply connected 5-manifold is diffeomorphic to a connected sum of manifolds that are listed in Table 1. Such a 5-manifold admits a totally real embedding into complex 5-space if and only if it is diffeomorphic to*

$$S^5\#\delta(S^3\widetilde{\times}S^2)\#(k_1-1)(S^2\times S^3)\#(k_2-1)(\mathrm{SU}(3)/\mathrm{SO}(3))\#M\#N \tag{3.5}$$

for

$$\delta + k_1 + k_2 + -1 = 0 \tag{3.6}$$

where $\delta \in \{0,1\}$, $k_1, k_2 \in \mathbb{N}$, the manifold M is a connected sum of an arbitrary number of copies of M_p^k and N is a connected sum of an arbitrary number of copies of the manifold X_k of Table 1. If the second Stiefel-Whitney class is zero, the explicit diffeomorphism type is given in (2.8).

3.3. **Proof of Corollary D.** A 5-manifold that satisfies the hypothesis of the corollary is diffeomorphic to S^5 or to a connected sum of copies of $S^2 \times S^3$ if its second Stiefel-Whitney class is zero and to a connected sum of $S^3\widetilde{\times}S^2$ with copies of $S^2 \times S^3$ if its second Stiefel-Whitney class is not zero according to Barden and Smale aforementioned classification results. Using a Mayer-Vietoris sequence and the universal coefficient theorem, it is immediate to compute

$$\hat{\chi}_{\mathbb{Z}/2}(S^3\widetilde{\times}S^2\#(n-1)(S^2\times S^3)) = n+1 \bmod 2 \tag{3.7}$$

and

$$\hat{\chi}_{\mathbb{Z}/2}(n(S^2 \times S^3)) = n + 1 \bmod 2 \tag{3.8}$$

for $n \in \mathbb{N}$. Item (B) of Theorem 2 implies that a sufficient and necessary condition for the totally real embedding to exists is for n to be an odd natural number.

3.4. **Proof of Theorem E.** Set $M := M^6$ and suppose there exists such a totally real immersion into $\mathbb{C}^6$. The triviality of the complexified tangent bundle $\mathbb{C}TM$ implies that $c_2(\mathbb{C}TM) = 0$. By definition of the Pontrjagin classes

$$p_i(M) = c_{2i}(\mathbb{C}TM), \tag{3.9}$$

it follows that $p_1(M) = 0$ if and only if $c_2(\mathbb{C}TM) = 0$. A 6-manifold that satisfies the hypothesis of Theorem E admits an almost-complex structure [19, Proposition 8], (cf. [23, Section 7]). The complexified tangent bundle of an almost-complex manifold M has the canonical eigenspaces decomposition

$$\mathbb{C}TM = T^{1,0}M \oplus T^{0,1}M, \tag{3.10}$$

into the holomorphic $T^{1,0}M$ and antiholomorphic $T^{0,1}M$ tangent bundles of M. The Chern classes of these bundles satisfy the equality

$$c_i(T^{1,0}M) = (-1)^i c_i(T^{0,1}M). \tag{3.11}$$

Suppose now that $p_1(M) = 0$. To prove the converse, we claim

$$c_i(\mathbb{C}TM) = 0 \tag{3.12}$$

for $i \in \mathbb{N}$; it then readily follows that $\mathbb{C}TM$ is trivial. From (3.9), we see that our hypothesis implies $c_2(\mathbb{C}TM) = 0$. Let us now show that $c_1(\mathbb{C}TM)$ is zero. It follows from using the Whitney product formula for Chern classes and (3.10) that

$$c_1(\mathbb{C}TM) = c_1(T^{1,0}M \oplus T^{0,1}M) = c_1(T^{1,0}M) + c_1(T^{0,1}M). \tag{3.13}$$

Identity (3.11) implies $c_1(\mathbb{C}TM) = 0$. Similarly, the Whitney product formula and identity (3.11) imply

$$c_3(\mathbb{C}TM) = c_1(T^{1,0}M) \cup c_2(T^{0,1}M) + c_2(T^{1,0}M) \cup c_1(T^{0,1}M) = 0. \tag{3.14}$$

Since $c_i(\mathbb{C}TM) = 0$ for every $i > 3$ due to dimensional reasons, we conclude that (3.12) holds. It follows that $\mathbb{C}TM$ is trivial and Theorem 1 implies the existence of the totally real immersion (1.9) as claimed.

The claims about the existence of a totally real embedding now follow from the first part of the theorem and Item (A) of Theorem 2.

□

3.5. **Proof of Theorem F: examples with arbitrary fundamental group.** Classical results of Dehn [4] and Kevaire-Milnor [18] imply that for any $N \geq 4$ and any finitely presented group G there exists a closed smooth stably parallelizable N-manifold $M^N(G)$ such that the fundamental group $\pi_1(M^N(G))$ is isomorphic to G. In particular, the complexified tangent bundle $\mathbb{C}TM^N(G)$ is trivial and a result of Gromov [7] implies that there is a totally real immersion $M^N(G) \to \mathbb{C}^N$ for every $N \geq 4$ as it is stated in Theorem 1. We now proceed to show the existence of a totally real embedding into $\mathbb{C}^N$. For the values $N = 4 + 2(k_1 - 1) + 3k_2$ with $k_1, k_2 \in \mathbb{N}$ the claim follows immediately by invoking Proposition 1 on the product of the 4-manifold $M^4(G)$ with $(k_1 - 1)$ copies of S^2 and k_2 copies of S^3. We apply Theorem 2 to deal with the cases N even and $N = 4k + 1$. A

standard computation using a Mayer-Vietoris sequence and the universal coefficient theorem implies that either $\hat{\chi}_{\mathbb{Z}/2}(M^5(G)) = 0$ or $\hat{\chi}_{\mathbb{Z}/2}(M^5(G)\#S^2 \times S^3) = 0$. Since $\pi_1(M^5(G)) = \pi_1(M^5(G)\#S^2 \times S^3)$, we conclude that the claim for $N = 5$ holds. Using product manifolds as before, we conclude that the claim regarding the existence of a totally real embedding holds for odd N. For even values of N, the argument is similar. A Mayer-Vietoris sequence reveals that the Euler characteristic of $M^N(G)$ is a positive even number. Taking connected sums of $M^N(G)$ with copies of $S^3 \times S^{N-3}$, one obtains a manifold with fundamental group G and zero Euler characteristic, which we continue to call $M^N(G)$.

□

Remark 3. *Johnson-Walton* [12, Theorem A] *pointed out that work of Kervaire* [13, 14] *implies that the manifolds of Theorem F are parallelizable.*

4. Examples of generic immersions

In this last section, we mention some examples of the following kind of immersions.

Definition 2. Let n be a nonnegative integer and $k \in \mathbb{N}$. An immersion

$$\pi : M^{2n+k} \to \mathbb{C}^{n+k} \tag{4.1}$$

is said to be generic if at each point $p \in M$ the real vector space

$$\pi_* TM \cap J\pi_* M \tag{4.2}$$

has dimension $2n$.

Definition 2 recovers the notion of totally real immersions of Definition 1 for the values $(n, k) = (0, N)$. A result of Hirsch [9] states that the bundle $TM^{2n+1} \oplus \epsilon$ is trivial if and only if M^{2n+1} immerses in $\mathbb{R}^{2n+2}$. Since every real hypersurface in $\mathbb{C}^{n+1}$ is automatically generic, any closed stably parallelizable $(2n+1)$-manifold has a generic immersion into $\mathbb{C}^{n+1}$. (cf. [10, Remark 4]). This yields the following examples of generic immersions.

Corollary 2. *A closed simply connected 5-manifold has a generic immersion into $\mathbb{C}^3$ if and only if its second Stiefel-Whitney class is zero, i.e., if the 5-manifold is stably parallelizable.*

For every $n \geq 2$ and every finitely presented group G there is a generic immersion

$$M^{2n+1}(G) \to \mathbb{C}^{n+1} \tag{4.3}$$

of a closed smooth orientable $(2n+1)$-manifold $M^{2n+1}(G)$ with fundamental group isomorphic to G.

References

[1] P. Ahern and W. Rudin, *Totally real embeddings of S^3 in $\mathbb{C}^3$*, Proc. Amer. Math. Soc., 94 (1985), 460 - 462.

[2] M. Audin, *Fibré normaux d'immersions en dimension double, points double d'immersions lagrangiennes et plongements totalement réels*, Comment. Math. Helvetici, 63 (1998), 593 - 623.

[3] D. Barden, *Simply connected five-manifolds*, Ann. of Math. 82 (1965), 365 - 385.

[4] M. Dehn, *Über unendliche diskontinuierliche Gruppen*, Math. Annalen 71 (1912), 16 - 144.

[5] F. Forstnerič, *On totally real embeddings into $\mathbb{C}^n$*, Expositiones Mathematicae, 4 (1986), 243 - 255.

[6] X. Gong, *On totally real spheres in complex space*, Math. Ann. 309 (1997), 611 - 623.

[7] M. Gromov, *Convex integration of differential relations I*, Math. USSR, Izv. 7 (1973), 329 - 343.
[8] M. Gromov, *Partial differential relations*, Ergebnisse der Mathematik und ihrer Grenzgebiete 9, Springer-Verlag, Berlin, 1986 x + 363 pp.
[9] M. W. Hirsch, *Immersions of manifolds*, Trans. Amer. Math. Soc. 93 (1959), 242 - 276.
[10] H. Jocobowitz and P. Landweber, *Manifolds admitting generic immersions into* $\mathbb{C}^N$, Asian J. Math. 11 (2007), 151 - 166.
[11] H. Jocobowitz and P. Landweber, *Totally real mappings and independent mappings*, Bull. Inst. Math. Acad. Sin. (N. S.) 8 (2013), 219 - 230
[12] F. E. A. Johnson and J. P. Walton, *Parallelizable manifolds and the fundamental group*, Mathematika 47 (2000), 165 - 172.
[13] M. A. Kervaire, *Relative characteristic classes*, Amer. J. Math. 79 (1957), 517 - 558.
[14] M. A. Kervaire, *Some non-stable homotopy groups of Lie groups*, Illinois J. Math. 4 (1960), 161 - 169.
[15] M. A. Kervaire, *Smooth homology spheres and their fundamental groups*, Trans. Amer. Math. Soc. 144 (1969), 67 - 72.
[16] J. Milnor, *A procedure for killing homotopy groups of differentiable manifolds*, Proc. Sympos. Pure Math., Vol. III, Amer. Math. Soc., Providence, R. I., 1961, pp. 39 - 55.
[17] G. Lusztig, J. Milnor, and F. Peterson, *Semi-characteristics and cobordism*, Topology 8 (1969), 357 - 359.
[18] J. Milnor and M. A. Kervaire, *Groups of homotopy spheres: I*, Ann. of Math. 77 (1963), 504 - 537.
[19] C. Okonek and A. Van de Ven, *Cubic forms and complex 3-folds*, L'Enseignement Mathématique 41 (1995), 297 - 333.
[20] R. H. Szczarba, *On tangent bundles of fibre spaces and quotient spaces*, Amer. J. Math. 86 (1964), 685 - 697.
[21] S. Smale, *On the structure of 5-manifolds*, Ann. of Math. 75 Vol. 2 (1962), 38 - 46.
[22] E. L. Stout and W. Zame, *A Stein manifold topologically but not holomorphically equivalent to a domain in* $\mathbb{C}^N$, Adv. Math. 60 (1986), 154 - 160.
[23] C. T. C. Wall, *Classification problems in differential topology V: On certain 6-manifolds*, Invent. Math. 1 (1966), 355 - 374.
[24] R. O. Wells, *Compact real submanifolds of a complex manifold with non-degenerate holomorphic tangent bundles*, Math. Ann. 179 (1969), 123 - 129.

Faculty of Education, University of Ljubljana, Kardeljeva Poščad 16, 1000, Ljubljana, Slovenia

Institute of Mathematics, Physics and Mechanics, Jadranksa 19, 1000, Ljubljana, Slovenia
E-mail address: marko.slapar@pef.uni-lj.si

Scuola Internazionale Superiori di Studi Avanzati (SISSA), Via Bonomea 265, 34136, Trieste, Italy
E-mail address: rtorres@sissa.it